

Data on Neutron Widths do not refute Random–Matrix Theory

J. F. Shriner^a, Jr., H. A. Weidenmüller^b, and G. E. Mitchell^{c*}

^a*Department of Physics, Tennessee Technological University, Cookeville, TN 38505, USA*

^b*Max-Planck-Institut für Kernphysik, 69029 Heidelberg, Germany*

^c*North Carolina State University, Raleigh, North Carolina 27695, USA and
Triangle Universities Nuclear Laboratory, Durham, North Carolina 27708, USA*

Recent analyses of the distribution of reduced neutron widths in the Nuclear Data Ensemble (NDE) and in the Pt isotopes claim strong disagreement with predictions of random–matrix theory. Using numerical simulations we identify the causes for the apparent disagreement. For the NDE it is a strong bias inherent in the method of analysis. In the Pt isotopes, it is likely to be a threshold state.

PACS numbers: 24.60.Ky, 24.60.Lz, 25.40.Ny, 29.87.+g

Recent analyses of the distribution of reduced neutron widths in the Pt isotopes and in the Nuclear Data Ensemble (NDE) have cast serious doubt on the validity of random–matrix theory (RMT) in nuclei. For the Pt isotopes, the analysis rejects agreement with RMT with a statistical significance of at least 99.997 per cent probability [1]. For the NDE, the corresponding figure is 99.17 per cent [2]. These results have found wide attention [3–6] eroding, as they seemingly do, one of the cornerstones of the statistical theory of nuclear reactions [7, 8]. In this Letter we show that the method of analysis used in Ref. [2] introduces a strong bias. The bias is responsible for the claimed disagreement of the NDE with RMT. In the Pt isotopes, a single–particle threshold state is probably responsible for the disagreement.

We briefly describe the method of analysis used in Ref. [2]. 1245 reduced neutron widths in the NDE of Refs. [9, 10] for 24 nuclides each containing between 14 and 178 resonances were used. A cutoff procedure [11] was used to eliminate p –wave resonances. The cutoff is based on the fact that s –wave (p –wave) resonance widths have an intrinsic energy dependence $E^{1/2}$ ($E^{3/2}$, respectively). Here E is the energy counted from neutron threshold, and the factors $E^{1/2}$ and $E^{3/2}$ have to be taken at the resonance energy. The transition to reduced widths removes the $E^{1/2}$ dependence of s –wave resonance widths. The remaining linear energy dependence of p –wave resonance widths was used to suppress these by introducing a cutoff that depends linearly on resonance energy and removing all widths smaller than the cutoff. The distribution of the remaining widths was analyzed using a maximum likelihood (ML) method. The method tests for agreement with the Porter–Thomas distribution (PTD) predicted by RMT. The PTD is a χ^2 –distribution with a single degree of freedom ($\nu = 1$). For comparison, distribution functions with other values of ν were admitted. The test determined which of these gave best agreement with the data. This led to the above–mentioned rejection of RMT.

The normalized χ^2 –distribution for ν degrees of freedom is

$$g(x, \nu, \langle x \rangle) = \frac{\nu}{2\Gamma(\nu/2)\langle x \rangle} \left(\frac{\nu x}{2\langle x \rangle} \right)^{(\nu/2)-1} \exp \left\{ -\frac{\nu x}{2\langle x \rangle} \right\}. \quad (1)$$

Here $\langle x \rangle$ denotes the average of x , and Γ is the complete gamma function. When the distribution is cut off at $x_{\min} > 0$ the resulting normalized function g_{cut} has the form $g_{\text{cut}}(x, \nu, \langle x \rangle) = (1/C)g(x, \nu, \langle x \rangle)\Theta(x - x_{\min})$ where Θ is the Heaviside function. With $T = x_{\min}/\langle x \rangle$ the normalization constant C is given by

$$C = \frac{\Gamma(\nu/2, \nu T/2)}{\Gamma(\nu/2)} \quad (2)$$

where $\Gamma(x, y)$ is the incomplete gamma function. Suppose now that for some fixed values of ν and $\langle x \rangle$, $g_{\text{cut}}(x, \nu, \langle x \rangle)$ is the correct distribution function for the variable x after cutoff. Then the normalized probability for x to have the measured value x_i at a resonance with energy E_i is $(1/C_i)g(x_i, \nu, \langle x \rangle)\Theta(x_i - x_{\min}(i))$. Here $x_{\min}(i)$ is the value of the cutoff at E_i , and C_i is given by Eq. (2), with $T = x_{\min}(i)/\langle x \rangle$. For a set of N_{incl} reduced widths x_i of resonances with energies E_i , $i = 1, \dots, N_{\text{incl}}$ remaining after the cutoff (so that $\Theta(x_i - x_{\min}(i)) = 1$ for all $i = 1, \dots, N_{\text{incl}}$) and included in the ML analysis, the likelihood function L is the product of the probabilities for finding resonance width x_i at energy E_i ,

$$L(\nu, \langle x \rangle) = \prod_{i=1}^{N_{\text{incl}}} \frac{1}{C_i} g(x_i, \nu, \langle x \rangle). \quad (3)$$

Now the argument is turned around: One considers L as a function of the two variables ν and $\langle x \rangle$ and argues that the maximum of L determines the most likely values of these parameters. The functional dependence of L on ν and $\langle x \rangle$ in the vicinity of the maximum determines the error and, thus, the statistical significance with which a value of ν that differs from the value at the maximum, is rejected. That is how the figure of 99.17 per cent cited above for the NDE is arrived at. In actual fact the method used in Ref. [2] is slightly different. Instead of the set of normalization factors C_i , a single joint

*Electronic address: jshriner@ntech.edu, Hans.Weidenmueller@mpi-hd.mpg.de

normalization factor was used. We return to that point below.

We have tested the method by simulation. Using a random-number generator we have drawn N real numbers y_i , $i = 1, \dots, N$ from a Gaussian probability distribution with unit width. The squares determine N widths $x_i = y_i^2$. In the limit $N \rightarrow \infty$, these by construction obey the PTD with $\langle x \rangle = 1$. (We have also simulated widths for $\langle x \rangle = 0.1$ and for $\langle x \rangle = 10$, with no discernible differences in the analysis.) Each of these widths is associated with one of N resonance energies $E_i = id$ with constant level spacing d [12]. We repeat that procedure 2500 times, thereby generating for each value of N an ensemble of widths and associated energies. For every member of the ensemble, we use the cutoff and apply the ML analysis described above. For clarity we distinguish the input value $\langle x \rangle = 1$ used to generate the data from the result of the analysis. The maxima of the 2500 likelihood functions L are located at values of ν denoted by $\hat{\nu}_k$ and of the average width denoted by $\langle \Gamma_n^0 \rangle_k$, $k = 1, \dots, 2500$. The uncertainty σ_k of $\hat{\nu}_k$ was estimated by determining those values of ν where $\ln L$ has decreased by the factor $1/2$. Since L is slightly asymmetric, σ_k was taken as the average of the two values. We calculated the mean values $\bar{\nu}$ of the $\hat{\nu}_k$ and $\langle \Gamma_n^0 \rangle$ of the widths. We determined the statistical error σ_{ens} of the ensemble average $\bar{\nu}$ from $\sigma_{\text{ens}}^{-2} = \sum_k \sigma_k^{-2}$.

A cutoff on the widths is defined by elimination of all normalized widths that are smaller than the cutoff value x_{min} . The normalized widths are the actual simulated widths x_i divided by their average. For finite N that average differs slightly from unity. The ML analysis is then applied to the remaining (unnormalized) $N_{\text{incl}} < N$ widths and the associated remaining energies. The latter do not have constant spacing. The likelihood function is given by Eq. (3), with $\langle x \rangle$ replaced by the unknown width $\langle \Gamma_n^0 \rangle$, with C_i given by Eq. (2), and with $T = x_{\text{min}}/\langle \Gamma_n^0 \rangle$. We have used six different cutoff procedures. (i) No cutoff ($x_{\text{min}} = 0$). (ii) A constant cutoff x_{min} independent of resonance energy as in Ref. [11]. Then all C_i have the same value (2). (iii) A linear cutoff $x_{\text{min}}(E_i) = X_{\text{min}}E_i/E_{\text{max}}$ where $E_{\text{max}} = Nd$ and where X_{min} measures the slope. With $T(E_i) = x_{\text{min}}(E_i)/\langle \Gamma_n^0 \rangle$ the normalization constants are $C_i = \Gamma(\nu/2, \nu T(E_i)/2)/\Gamma(\nu/2)$. (iv) A linear cutoff as under (iii) but for a modified form of the likelihood function (3) where every C_i is replaced by C , the linear smooth average of the C_i ,

$$C = \frac{1}{E_{\text{max}}} \int_0^{E_{\text{max}}} dE \frac{\Gamma(\nu/2, \nu T(E)/2)}{\Gamma(\nu/2)}, \quad (4)$$

and $E_{\text{max}} = Nd$. That is the form [13] of L actually used in Refs. [1, 2]. (v) A cutoff given by $x_{\text{min}}(E_i) = X_{\text{min}}[E_i/E_{\text{max}}]^{3/2}$. That is the cutoff used in Ref. [1] for the reduced neutron widths in the Pt isotopes. The normalization constants C_i have the same form as under (iii). (vi) Same as under (v) but with every C_i replaced by C as defined in Eq. (4). Method (vi) was used in Ref. [1]. Method (iii) (method (v)) corresponds to the

standard ML approach and is supported by mathematical theory while method (iv) (method (vi), respectively) is not. We test whether methods (iv) or (vi) yield results that differ from those of methods (iii) or (v), respectively.

We have simulated data for $N = 100, N = 450, N = 1100$. Here $N = 100$ corresponds roughly to a typical set of widths for a given nuclide in the NDE, $N = 1100$ is approximately the total number of widths in the NDE, and $N = 450$ provides an intermediate value. The three values of N are also used to test whether there is a systematic N -dependent trend in the ML analysis. Some of our results are shown in the tables. In Table I we show for methods (i) to (iv) and the two extreme values of N the cutoff parameters as defined above, the average number $\bar{N}_{\text{incl}}$ of widths retained after cutoff, and the ensemble averages $\bar{\nu}$ and $\langle \Gamma_n^0 \rangle$ with their statistical errors resulting from ensemble averaging. Since we used 2500 simulations, the errors are small in all cases. Method (i) serves as a test of our method. Method (i) yields for all three N -values $\bar{\nu} \approx 1$ and $\langle \Gamma_n^0 \rangle \approx 1$, consistent with the input. The values of $\bar{\nu}$ and $\langle \Gamma_n^0 \rangle$ determined with methods (ii) to (iv) increase with increasing cutoff and decrease very slowly with increasing N . Methods (iii) and (iv) give very similar results for all three values of N ; the symmetrization (4) does not adversely affect the results. In view of the small statistical error, agreement with RMT would be firmly excluded for all non-zero cutoffs that are bigger than 0.02 – in spite of the fact that the data have been generated from the PTD. These facts establish the existence of a strong cutoff-dependent bias in the ML analysis of neutron resonance widths [14].

How does the bias arise? How can it be that the PTD used to generate the data does not give the best fit to these same data? Let us consider method (ii) (constant cutoff at x_{min}). An unbiased analysis would test whether after removal of all widths smaller than x_{min} , the shape of the distribution of the remaining widths is that of a PTD with a cutoff at x_{min} . The analysis might, for instance, test for maximum overlap, irrespective of normalization. The ML analysis, in contradistinction, assigns a normalized probability to each width and searches for the maximum of the product of these probabilities. That introduces the factor $C^{-N_{\text{incl}}} = [\Gamma(\nu/2, \nu T/2)/\Gamma(\nu/2)]^{-N_{\text{incl}}}$ in Eq. (3). That factor has a non-trivial dependence on ν and, through T , on $\langle \Gamma_n^0 \rangle$. The maximum of L is, thus, not determined by maximum overlap. It is influenced by the normalization factors. That causes the bias. Similar considerations apply for non-constant cutoffs (methods (iii) to (vi)).

The statistical error on $\bar{\nu}$ is small and would decrease further if the number of simulations were increased. However, the maximum $\hat{\nu}_k$ of L for an individual simulation with index k does not, in general, coincide with $\bar{\nu}$. For the cases considered in Table I, Table II shows the root-mean-square deviations $\delta\nu$ of the $\hat{\nu}_k$ from $\bar{\nu}$. Such deviations are likewise expected to occur for a set of N widths in any given nucleus. Table II also shows the average $\bar{\sigma}$ of the widths σ_k defined above. We note that $\bar{\sigma} \geq \delta\nu$ in

TABLE I: Ensemble averages $\overline{\nu}$ and $\overline{\langle \Gamma_n^0 \rangle}$ for different sample sizes, different methods, and different cutoff parameters.

N	$x_{\min}/X_{\min}^b$	$\overline{N_{\text{incl}}}^c$	Method (ii)		$\overline{N_{\text{incl}}}^{ac}$	Method (iii)		Method (iv) ^a	
			$\overline{\nu}$	$\overline{\langle \Gamma_n^0 \rangle}$		$\overline{\nu}$	$\overline{\langle \Gamma_n^0 \rangle}$	$\overline{\nu}$	$\overline{\langle \Gamma_n^0 \rangle}$
100	0 ^d	100	0.994 ± 0.002	0.963 ± 0.003		Same as Method (ii)		Same as Method (ii)	
	0.01	92	0.997 ± 0.004	1.043 ± 0.003	95	0.999 ± 0.004	1.016 ± 0.003	1.002 ± 0.004	1.016 ± 0.003
	0.02	89	1.003 ± 0.004	1.079 ± 0.003	93	0.999 ± 0.004	1.039 ± 0.003	1.003 ± 0.004	1.039 ± 0.003
	0.05	83	1.034 ± 0.005	1.158 ± 0.004	88	1.003 ± 0.004	1.087 ± 0.003	1.009 ± 0.004	1.087 ± 0.003
	0.10	75	1.111 ± 0.006	1.258 ± 0.004	83	1.024 ± 0.005	1.146 ± 0.004	1.031 ± 0.005	1.146 ± 0.004
	0.20	66	1.303 ± 0.008	1.419 ± 0.004	77	1.081 ± 0.006	1.236 ± 0.004	1.089 ± 0.006	1.236 ± 0.004
	0.30	58	1.532 ± 0.009	1.560 ± 0.005	72	1.141 ± 0.006	1.311 ± 0.004	1.149 ± 0.006	1.310 ± 0.004
1100	0 ^d	1100	0.9999 ± 0.0007	0.9965 ± 0.0008		Same as Method (ii)		Same as Method (ii)	
	0.01	1012	1.009 ± 0.001	1.082 ± 0.001	1042	1.003 ± 0.001	1.052 ± 0.001	1.003 ± 0.001	1.052 ± 0.001
	0.02	977	1.018 ± 0.001	1.121 ± 0.001	1017	1.007 ± 0.001	1.077 ± 0.001	1.007 ± 0.001	1.077 ± 0.001
	0.05	906	1.056 ± 0.002	1.206 ± 0.001	971	1.018 ± 0.001	1.128 ± 0.001	1.018 ± 0.001	1.128 ± 0.001
	0.10	828	1.137 ± 0.002	1.313 ± 0.001	917	1.041 ± 0.002	1.190 ± 0.001	1.041 ± 0.002	1.190 ± 0.001
	0.20	721	1.327 ± 0.002	1.485 ± 0.001	844	1.100 ± 0.002	1.287 ± 0.001	1.099 ± 0.002	1.287 ± 0.001
	0.30	643	1.548 ± 0.003	1.635 ± 0.002	789	1.162 ± 0.002	1.365 ± 0.001	1.159 ± 0.002	1.365 ± 0.001

^a $\overline{N_{\text{incl}}}$ is always the same for Methods (iii) and (iv).

^bFor method (ii), $x_{\min}$ is the relevant cutoff parameter; for methods (iii) and (iv), $X_{\min}$ is the quantity that must be specified.

^cUncertainties in $\overline{N_{\text{incl}}}$ are much less than 1.

^d $x_{\min} = 0$ or $X_{\min} = 0$ corresponds to method (i).

TABLE II: Error estimates for $N = 100$ and $N = 1100$ for individual spectra

N	$x_{\min}/X_{\min}^a$	Method (ii)		Methods (iii)-(iv)	
		$\delta\nu$	$\overline{\sigma}$	$\delta\nu$	$\overline{\sigma}$
100	0 ^b	0.13	0.12	Same as Method (ii)	
	0.01	0.20	0.20	0.15	0.18
	0.02	0.21	0.23	0.15	0.20
	0.05	0.24	0.27	0.16	0.22
	0.10	0.25	0.32	0.17	0.25
	0.20	0.28	0.40	0.19	0.29
	0.30	0.31	0.47	0.20	0.32
1100	0 ^b	0.03	0.03	Same as Method (ii)	
	0.01	0.06	0.06	0.05	0.05
	0.02	0.06	0.07	0.06	0.06
	0.05	0.07	0.08	0.06	0.07
	0.10	0.07	0.09	0.07	0.07
	0.20	0.08	0.11	0.07	0.08
	0.30	0.09	0.13	0.08	0.09

^aFor method (ii), $x_{\min}$ is the relevant cutoff parameter; for methods (iii) and (iv), $X_{\min}$ is the quantity that must be specified.

^b $x_{\min} = 0$ or $X_{\min} = 0$ is the same for each method; these are the results when method (i) is utilized.

all cases where a cutoff is utilized. If all σ_k were equal, the statistical error on $\overline{\nu}$ in Table I would simply be given by $\overline{\sigma}/\sqrt{K}$ with $K = 2500$.

In Ref. [2] the weighted average of $\hat{\nu}_k$ over 24 nuclides is

given as $\overline{\nu}_{\text{Koe}} = 1.217 \pm 0.092$. Because of the small number of nuclides and because of the spread of the number of resonances per nuclide, the error on $\overline{\nu}_{\text{Koe}}$ is much larger than for the 2500 simulations in our Table I. Within that error, the result for $\overline{\nu}_{\text{Koe}}$ agrees, however, with the results of our Table I for cutoff values bigger than 0.2. In other words, the NDE behaves exactly as is expected for data that follow the PTD. Thus, the conclusions drawn in Ref. [2] on the NDE cannot be upheld. The authors of Ref. [10] have concluded that the distribution of neutron widths in the NDE agrees with the PTD. There is no reason to doubt that conclusion.

The Pt isotopes are special. While the analysis of the NDE in Ref. [2] yields for all nuclides values for $\hat{\nu}_k$ that are bigger than unity or, within errors, are consistent with unity, the values for ν_{exp} given in Table I of Ref. [1] for the three Pt isotopes range from 0.47 to 0.60 with errors not larger than 0.28. While in Ref. [2] the NDE is claimed to disagree with RMT because $\overline{\nu}_{\text{Koe}}$ is too big, the disagreement claimed in Ref. [1] comes about because ν_{exp} is too small. That suggests that the distribution of widths in the Pt isotopes is significantly different from that of the nuclides in the NDE, and that a special mechanism may here be at work. The investigation of the Pt isotopes in Ref. [1] was motivated by the fact that for these mass numbers, the neutron strength function is close to its maximum, so that p -wave contributions are effectively suppressed by about a factor 10 [1]. The enhancement of the s -wave strength function is due to a threshold state: With increasing mass number, the $4s$ single-particle state of the shell model is pulled into the

TABLE III: Values of $\overline{\nu}$ for spectra with $N = 161$ and $E_{\max} = 15.93$ keV for different values of $|E_0|$ when method (vi) is applied.

$X_{\min}$	$ E_0 $ (keV)		
	0.5	5	20
0	0.797 ± 0.001	0.890 ± 0.002	0.970 ± 0.002
0.01	0.690 ± 0.002	0.851 ± 0.002	0.976 ± 0.003
0.02	0.676 ± 0.002	0.846 ± 0.003	0.979 ± 0.003
0.05	0.666 ± 0.002	0.850 ± 0.003	0.992 ± 0.003
0.10	0.674 ± 0.003	0.866 ± 0.003	1.011 ± 0.004
0.20	0.704 ± 0.003	0.908 ± 0.004	1.057 ± 0.004
0.30	0.733 ± 0.003	0.946 ± 0.004	1.098 ± 0.004

nuclear potential well, and it passes threshold at or near the masses of the Pt isotopes. In Ref. [4] it was shown that a threshold state at energy $|E_0|$ either above or below threshold modifies the s -wave neutron widths by the factor

$$f^2(E) = \frac{\sqrt{E}}{E + |E_0|}. \quad (5)$$

Here $\sqrt{E}$ is the usual s -wave penetration factor, and $1/(E + |E_0|)$ is the resonance enhancement due to the threshold state. For each resonance, $f^2(E)$ has to be taken at the resonance energy E_i . In Ref. [1], the reduced widths of the Pt isotopes were analyzed using an energy-dependent cutoff proportional to $E^{3/2}$ without taking into account the denominator in Eq. (5). To test for the influence of the threshold state, we simulated 2500 sets of N widths x_i and N equally spaced energies E_i in the interval $\{0, E_{\max}\}$ as described above, with $N = 153$ and $E_{\max} = 4.98$ keV for ^{192}Pt and $N = 161$ and $E_{\max} = 15.93$ keV for ^{194}Pt , plus some other cases for additional information. For each realization of the ensemble, we multiplied every x_i by $1/(E_i + |E_0|)$. (Other energy-independent factors contributing to f^2 were ignored since a rescaling of all widths does not affect the value of $\hat{\nu}_k$). The resulting set of modified widths and associated resonance energies was subject to the ML analysis with an energy-dependent cutoff.

Some of our results are shown in Table III. Without

cutoff, significant deviations from the input value $\nu = 1$ appear only when $|E_0|/E_{\max} \leq 1/2$ or so. For the values of $|E_0|$ shown, these deviations are considerably smaller than the experimental ones in the Pt isotopes. Methods (iii) and (iv) again give very similar results; for some combination of values of $|E_0|$ and cutoff parameter, values of $\overline{\nu}$ as small as 0.7 are found. The results of methods (v) and (vi) differ, method (v) generally giving smaller values for $\overline{\nu}$. In both cases these values first drop and then increase with increasing cutoff, the values for method (vi) attaining a minimum around 0.70 for $|E_0| = 0.5$ keV. That is close to but larger than the experimental values that range from 0.5 to 0.6. However, the replacement $x_i \rightarrow f^2(E_i)x_i$ is only approximately valid if $|E_0|$ is as small as 1 keV or so. It is shown in Ref. [4] that in such a case, the entire R -matrix analysis of the scattering data must be redone, taking the existence of the threshold state into account from the outset. It is not known how exactly that affects the distribution of the x_i . We conclude that a single-particle state close to threshold is a likely candidate for the deviations found in the Pt isotopes. Establishing its existence would require, however, a complete re-analysis of the scattering data that would not make use of standard R -matrix codes.

In summary, we have shown that the ML analysis used in Ref. [2] produces a strong bias. The bias distorts the value of $\overline{\nu}$ determined from the data and leads to an apparent severe disagreement with RMT predictions. When the bias is taken into account, the distribution of neutron widths in the NDE is consistent with RMT. For the Pt isotopes we have shown that a single-particle threshold state is a likely cause for the observed disagreement with RMT. To firmly establish this assertion, a complete re-analysis of the scattering data would be required, however.

Acknowledgments

One of us (JFS) thanks P. E. Koehler for very helpful discussions concerning his application of the maximum likelihood method. One of us (GEM) acknowledges the support of the US Department of Energy via grants No. DE-FG52-09NA29460 and No. DE-FG02-97-ER41042

-
- [1] P. E. Koehler, F. Becvar, M. Krticka, J. A. Harvey, and K. H. Guber, Phys. Rev. Lett. **105**, 072502 (2010).
 - [2] P. E. Koehler, Phys. Rev. C **84**, 034312 (2011).
 - [3] E. S. Reich, Nature **466**, 1034 (2010).
 - [4] H. A. Weidenmüller, Phys. Rev. Lett. **105**, 232501 (2010).
 - [5] G. L. Celardo et al., Phys. Rev. Lett. **106**, 042501 (2011).
 - [6] A. Volya, Phys. Rev. C **83**, 044312 (2011).
 - [7] H. A. Weidenmüller and G. E. Mitchell, Rev. Mod. Phys. **81**, 539 (2009).
 - [8] G. E. Mitchell, A. Richter, and H. A. Weidenmüller, Rev. Mod. Phys. **82**, 2845 (2010).
 - [9] R. U. Haq, A. Pandey, and O. Bohigas, Phys. Rev. Lett. **48**, 1086 (1982).
 - [10] O. Bohigas, R. U. Haq, and A. Pandey, in *Nuclear Data for Science and Technology*, edited by K. H. Böckhoff (D. Reidel, Dordrecht, 1983), p. 809.
 - [11] H. S. Camarda, Phys. Rev. C **49**, 1391 (1994).
 - [12] In choosing a constant level spacing d we neglect the small spacing fluctuations that actually occur in mea-

sured data. These are irrelevant for this analysis.

- [13] P. E. Koehler, private communications to J. F. Shriner, Jr. (2011-2012).
- [14] In contrast to Refs. [1, 2], the analysis of Ref. [11] did correct for the bias.